| AD | ) |  |
|----|---|--|
|    |   |  |

Award Number: DAMD17-98-1-8108

TITLE: National Hospital Discharge Survey Data Analysis of Breast Cancer Between African American and White Women

PRINCIPAL INVESTIGATOR: Kyung Sook Kim, Ph.D.

CONTRACTING ORGANIZATION: Howard University
Washington, DC 20059

REPORT DATE: August 2000

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

| •   | REPORT DOCUM  | MENTATION P  |  | Form Approved<br>MB No. 074-0188   |
|---|---|--|--|--|
| Public reporting burden for this collection of inform maintaining the data needed, and completing and reincluding suggestions for reducing this burden to WVA 22202-4302, and to the Office of Managemen                      | eviewing this collection of information. Send<br>Vashington Headquarters Services, Directorat   | d comments regarding this burden<br>te for Information Operations and  | ewing instructions, set estimate or any othe Reports 1215 leffer.  | arching existing data sources, gathering and   |
| 1. AGENCY USE ONLY (Leave blank)  | 2. REPORT DATE<br>August 2000   | 3. REPORT TYPE AND Annual Summary  | DATES COVERE   |  |
| 4. TITLE AND SUBTITLE National Hospital Discha Cancer Between African A   |   | ysis of Breast   | 5. FUNDING N<br>DAMD17-98  | NUMBERS  |
| 6. AUTHOR(S)  Kyung Sook Kim, Ph.D.   |   |  |  |  |
| 7. PERFORMING ORGANIZATION NAM<br>Howard University<br>Washington, DC 20059   | IE(S) AND ADDRESS(ES)   |  | 8. PERFORMIN<br>REPORT NU  | IG ORGANIZATION<br>IMBER   |
| E-MAIL:<br>kkim@fac.howard.edu  |   |  |  |  |
| 9. SPONSORING / MONITORING AGER<br>U.S. Army Medical Research and M<br>Fort Detrick, Maryland 21702-5012  | fateriel Command  | •)   |  | ING / MONITORING<br>REPORT NUMBER  |
| 11. SUPPLEMENTARY NOTES   |   |  |  |  |
| 12a. DISTRIBUTION / AVAILABILITY S'<br>Approved for public release; distribu  |   |  |  | 12b. DISTRIBUTION CODE   |
| 13. ABSTRACT (Maximum 200 Words)  | 1   |  |  | <u> </u>   |
| American (AA) and V (NHDS) data from 19 Specific standard error program. The statistic two-sided z-test with cases decreased amore associated with the local linearly decreasing ov Mean number of diag that of AA was not li | timated age-specific hospit White women through ana 988 to 1994. Weighted lir ors for rates analyzed in the ical significance of the weighted a critical value of 1.96 (and many many many many many many many many | alyzing National Host<br>near regression was<br>ne regression were p<br>ighted least squares<br>=0.05). Hospital act<br>of the least squares<br>of the least squares | spital Dischar<br>used to analy<br>roduced using<br>test for trend<br>dmission rate<br>A women we<br>he mean 'Ler<br>of for AA wo<br>mong whites | rge Survey yze trends. ng the SUDAAN ds is based on the es of breast cancer ere cubically ngth of stay' was omen (p<0.43). (p<0.005) while |
| 14. SUBJECT TERMS   |   | 1  | . 1  | 15. NUMBER OF PAGES  |
| Breast Cancer, National H<br>admission rate due to br   | -   | rvey data, Hospi   |  | 20<br>16. PRICE CODE   |

NSN 7540-01-280-5500

OF REPORT

17. SECURITY CLASSIFICATION

Unclassified

20. LIMITATION OF ABSTRACT

Unlimited

19. SECURITY CLASSIFICATION

Unclassified

OF ABSTRACT

18. SECURITY CLASSIFICATION

Unclassified

OF THIS PAGE

#### FOREWORD

| Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.   |
|---|
| Where copyrighted material is quoted, permission has been obtained to use such material.  |
| Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.  |
| Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.   |
| N/A In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985). |
| $\underline{X}$ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.  |
| N/A In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by   |

investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research utilizing recombinant DNA, the

the National Institutes of Health.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Kyungstoolli 8/22/00

Date

Date

## **Table of Contents**

| I   | oage |
|---|------|
| Cover   | . 1  |
| SF 298  | 2    |
| Foreword  | 3    |
| Table of Contents   | 4    |
| Introduction  | . 5  |
| Body  | . 5  |
| Conclusions   | . 7  |
| Appendices  | . 8  |
| Table 1. Hospital admission rates and 95% confidence interval of breast                                   |      |
| cancer between White and African American women from 1988 to 1994   | . 9  |
| Table 2. Age-specific admission rates of breast cancer among White women from 1988 to 1994                | . 10 |
| Table 3. Age-specific admission rates of breast cancer among African American women from 1988 to 1994     |      |
| Table 4. Distribution of breast cancer patients age-group by race from 1988 to                            | 12   |
| Table 5. Mean (s.e.) of Age, length of stay, number of diagnosis and procedures by race from 1988 to 1994 |      |
| Table 6. Distribution of Expected sources of payment (ESOP) by race from 1988 to 1994                     |      |
| Figure 1. Hospital admission rates and 95% C.I. of breast cancer by race from 1988 to 1994                |      |
| Figure 2. Hospital admitted breast cancer percentages (± s.e.) of ages 20-44 by race from 1988 to 1994    |      |
| Figure 3. Mean age (95% C.I.) of breast cancer patients by race from 1988 to 1994                         | 17   |
| Figure 4. Average length of stay (95% C.I.) of breast cancer patients by race from 1988 to 1994           |      |
| Figure 5. Mean number of diagnosis and procedures (95% C.I.) by race from 1988 to 1994                    |      |
| Figure 6. Percentages of routine discharge (±s.e.) by race from 1988 to 1994                              |      |

#### Introduction

The present study attempts to estimate age-specific hospital admission rates for breast cancer in African American (AA) and White women through analyzing National Hospital Discharge Survey (NHDS) data from 1988 to 1994 with 1987 data used for baseline comparison. Due to the sample design change in 1988, the comparison with 1987 as a baseline may not be valid nor meaningful. Therefore, this report will include only 1988 to 1994 data, excluding 1987 data.

Trend analyses on age-specific and overall breast cancer admissions were included in this report.

#### Body

We analyzed age-specific hospital admission rates for breast cancer from 1990 to 1994 by using NHDS data. In order to estimate standard errors of age-specific rates, we went to the Research Data Center for National Center for Health Statistics to use "restricted data" such as 'Hospital ID' and 'primary sampling unit'. Table 1 and Figure 1 show the hospital admission rates and 95% confidence interval of breast cancer between White and African American women from 1988 to 1994. Weighted linear regression was used to analyze trends. Specific standard errors for rates analyzed in the regression were produced using the SUDAAN program. The statistical significance of the weighted least squares test for trends is based on the two-sided z-test with a critical value of 1.96 ( $\alpha$ =0.05). Hospital admission rates of breast cancer were linearly decreasing among White women (p<0.0004). Among AA women those rates were not linearly associated with time (p<0.54). For AA women, the lowest rate occurred in 1990 and there was a little increase in 1993.

Table 2 and 3 show age-specific hospital admission rates of breast cancer by race from 1988 to 1994. Among White women, excluding age group 20-44, hospital admission rates

significantly decreased over the study period. Only ages 60-69 showed a significant linear decrease among AA women.

Table 4 shows percentages by age groups, who were admitted in the hospital for breast cancer. AA women ages 20-44 showed a significant increase over the study period (p<0.01). Figure 2 shows the percentages of women ages 20-44, who were hospitalized due to breast cancer over the study period by race. The trend analysis supported the hypothesis that AA women had higher proportion of breast cancer in the younger age group compared to White women.

Table 5 shows means and standard errors of age, length of stay, number of diagnosis, and procedures by race. As Figure 3 shows that hospital admitted breast cancer patients mean ages were stable among Whites, compared to AA women which decreased significantly over the study period (p<0.0001). Figure 4 shows "Length of stay" means over time by race. These means were linearly decreasing over time for Whites (p<0.0093) but not for AA (p<0.43). Figure 5 shows means of "number of diagnosis" and "number of procedures" over time. The mean number of diagnosis was significantly increasing over time among Whites (p<0.005) while that of AA was not linearly associated (p<0.24). The mean number of procedures was not linearly increasing among Whites (p<0.13) nor among AA women (p<0.13).

Figure 6 shows percentages of "routine discharge" by race over the study years. No significant trends exist for either race. Table 6 shows the percentages of expected sources of payments by year and race. Approximately 40% of the White women were on Medicare versus 30% of the AA women. This trend decrease among AA women over study period (p<0.05). AA women were five times (20%) more likely to be covered by Medicaid than Whites (4%). On average, 54% of breast cancer patients who were AA were from the Southern region and 5%

from the Western region, while 35% of the White from the South and 13% from the West.

### **Conclusions**

Hospital admission rates of breast cancer were linearly decreasing among White women. Those rates were cubically associated with the lowest rate in 1990 and an increase in 1993 among AA women. Hospital admission rates of breast cancer among women ages 20-44 did not show any trends over the study period. However, the percentage of hospital admitted White and AA women ages 20-44 support the hypothesis that AA women have a higher proportion of breast cancer among younger age groups compared to White women.

## **APPENDICES**

Table 1. Hospital admission rates\* and 95% confidence interval of breast cancer between White and African American women from 1988 to 1994.

|       | V        | /hite                   | Afri     | can American            |
|-------|----------|-------------------------|----------|-------------------------|
| Year  | Rate     | 95% Confidence Interval | Rate     | 95% Confidence Interval |
| 1988  | 310.95   | (262.12, 359.78)        | 396.82   | (251.40, 542.25)        |
| 1989  | 290.25   | (245.36, 335.14)        | 299.95   | (202.51, 397.40)        |
| 1990  | 265.47   | (232.14, 298.80)        | 182.47   | (120.45, 244.50)        |
| 1991  | 248.66   | (215.99, 281.33)        | 198.80   | (143.75, 253.86)        |
| 1992  | 261.23   | (224.23, 298.22)        | 228.37   | (138.07, 318.67)        |
| 1993  | 243.06   | (211.50, 274.62)        | 252.75   | (190.22, 315.27)        |
| 1994  | 217.11   | (186.13, 248.09)        | 219.40   | (153.75, 285.05)        |
| Trend | Z = -3.5 | 5339 P <0.0004          | Z = -0.6 | 6152 P < 0.54           |

rates\* per 100,000

Table 2. Age-specific admission rates\* of breast cancer among White women from 1988 to 1994

| Age group 20-44 |          | -44      | 45-59   |        | 60-6    | 60-69  |           | 1       |
|-----------------|----------|----------|---------|--------|---------|--------|-----------|---------|
| Year            | Rate     | s.e.     | Rate    | s.e.   | Rate    | s.e.   | Rate      | s.e.    |
| 1988            | 100.43   | 12.88    | 498.13  | 71.09  | 531.48  | 45.04  | 685.34    | 62.33   |
| 1989            | 83.29    | 10.14    | 467.04  | 59.16  | 533.37  | 52.84  | 638.40    | 62.66   |
| 1990            | 94.61    | 12.32    | 391.01  | 35.57  | 537.04  | 49.17  | 499.11    | 39.62   |
| 1991            | 84.99    | 10.46    | 322.69  | 30.61  | 500.89  | 52.79  | 563.18    | 51.13   |
| 1992            | 89.42    | 10.68    | 387.24  | 39.37  | 486.01  | 61.04  | 538.88    | 56.30   |
| 1993            | 97.43    | 15.48    | 301.43  | 28.53  | 433.07  | 46.19  | 550.58    | 50.69   |
| 1994            | 75.55    | 9.40     | 354.82  | 39.76  | 384.00  | 41.08  | 389.40    | 47.20   |
| Trend           | z = -1.0 | 5 p<0.29 | z=-2.58 | p<0.01 | z=-2.99 | p<0.03 | z = -3.31 | p<0.009 |

rates\* per 100,000

Table 3. Age-specific admission rates\* of breast cancer among African American women from 1988 to 1994.

| Age grou | up 20-4  | 14     | 45-5     | 9      | 60-69     |         | 70-84   |        |
|----------|----------|--------|----------|--------|-----------|---------|---------|--------|
| Year     | Rate     | s.e.   | Rate     | s.e.   | Rate      | s.e.    | Rate    | s.e.   |
| 1988     | 90.39    | 17.85  | 636.14   | 248.78 | 969.20    | 189.92  | 1410.99 | 616.46 |
| 1989     | 100.50   | 25.65  | 393.09   | 105.11 | 598.98    | 157.01  | 1204.00 | 412.15 |
| 1990     | 69.18    | 16.69  | 257.73   | 87.55  | 371.98    | 117.56  | 588.62  | 203.90 |
| 1991     | 75.97    | 17.04  | 433.34   | 78.72  | 425.42    | 134.78  | 306.89  | 77.62  |
| 1992     | 99.81    | 23.72  | 539.51   | 210.96 | 383.60    | 97.36   | 272.59  | 80.66  |
| 1993     | 102.85   | 25.17  | 483.91   | 93.25  | 404.56    | 112.84  | 625.64  | 180.90 |
| 1994     | 109.33   | 30.36  | 421.37   | 105.80 | 328.52    | 70.31   | 404.49  | 101.31 |
| Trend    | z = 0.60 | p<0.55 | z = 0.81 | p<0.42 | z = -2.52 | p< 0.12 | z=-0.40 | p<0.69 |

rates\* per 100,000

Table4. Distribution of breast cancer patients age-group by race from 1988 to 1994


|           |      | White     |          | African American      |
|-----------|------|-----------|----------|-----------------------|
| Age-group | Year | %         | s.e.     | % s.e.                |
|           | 1988 | 17.44     | 2.07     | 14.07 3.53            |
|           | 1989 | 15.47     | 1.35     | 20.69 4.86            |
|           | 1990 | 19.06     | 1.79     | 23.25 4.55            |
| 20-44     | 1991 | 18.39     | 1.74     | 23.85 4.99            |
|           | 1992 | 18.18     | 1.87     | 27.15 6.83            |
|           | 1993 | 21.11     | 2.63     | 25.14 5.17            |
|           | 1994 | 18.17     | 2.09     | 30.6 6.31             |
| Trend     |      | z = 1.38  | p < 0.17 | z = 2.63 p<0.01       |
|           | 1988 | 32.77     | 3.26     | 31.56 9.39            |
|           | 1989 | 33.09     | 2.39     | 25.85 6.14            |
|           | 1990 | 30.28     | 1.9      | 27.84 7.89            |
| 45-59     | 1991 | 26.64     | 1.64     | 41.75 4.87            |
|           | 1992 | 31.6      | 2.24     | 46.18 10.13           |
|           | 1993 | 27.2      | 1.95     | 38.28 6.29            |
|           | 1994 | 36.91     | 2.58     | 39.56 7.11            |
| Trend     |      | z = -0.01 | p<0.99   | z = 1.74 $p < 0.08$   |
|           | 1988 | 22.42     | 1.76     | 25.35 6.12            |
|           | 1989 | 23.93     | 1.89     | 20.73 5.72            |
|           | 1990 | 26.19     | 1.86     | 21 6.47               |
| 60-69     | 1991 | 25.57     | 2.14     | 21.34 4.98            |
|           | 1992 | 23.07     | 1.96     | 16.52 4.91            |
|           | 1993 | 21.66     | 1.98     | 15.58 3.67            |
|           | 1994 | 21.05     | 1.94     | 14.44 3.04            |
| Trend     |      | z= -1.07  | p < 0.29 | z = -1.95 $p < 0.051$ |
|           | 1988 | 27.36     | 2.25     | 29.01 9.69            |
|           | 1989 | 27.51     | 2.32     | 32.73 7.08            |
|           | 1990 | 24.47     | 1.71     | 27.91 7.25            |
| 70-84     | 1991 | 29.4      | 2.1      | 13.07 3.2             |
|           | 1992 | 27.14     | 2.36     | 10.15 3.31            |
|           | 1993 | 30.03     | 2.38     | 21.00 5.3             |
|           | 1994 | 23.87     | 2.22     | 15.41 3.79            |
| Trend     |      | z = -1.58 | p < 0.11 | z = -1.79 $p < 0.07$  |

Table 5. Mean (s.e.) of Age, length of stay, number of diagnosis and procedures by race from 1988 to 1994

|           |      | White       |            | African Aeı | rican      |  |
|-----------|------|-------------|------------|-------------|------------|--|
|           | Year | Mean        | s.e.       | Mean        | s.e.       |  |
|           | 1988 | 58.82       | 0.82       | 59.38       | 1.68       |  |
| Age       | 1989 | 59.3        | 0.67       | 59.23       | 2.46       |  |
| · ·       | 1990 | 58.26       | 0.64       | 56.87       | 1.99       |  |
|           | 1991 | 59.74       | 0.61       | 54.24       | 1.41       |  |
|           | 1992 | 58.64       | 0.66       | 52.41       | 1.57       |  |
|           | 1993 | 59.05       | 0.91       | 54.59       | 1.79       |  |
|           | 1994 | 58.01       | 0.69       | 52.01       | 1.44       |  |
| Trend     |      | z = -0.8036 | p= 0.4216  | z = -3.8503 | p = 0.0001 |  |
|           | 1988 | 5.99        | 0.4        | 6.04        | 0.9        |  |
|           | 1989 | 5.86        | 0.42       | 6.68        | 0.96       |  |
| Length    | 1990 | 4.83        | 0.23       | 8.34        | 1.03       |  |
| of stay   | 1991 | 5.63        | 0.4        | 6.43        | 0.96       |  |
|           | 1992 | 4.76        | 0.31       | 6.17        | 0.7        |  |
|           | 1993 | 5.05        | 0.45       | 5.89        | 0.6        |  |
|           | 1994 | 4.66        | 0.29       | 6.37        | 1.88       |  |
| Trend     |      | z = -2.6022 | p = 0.0093 | z = -0.7803 | p = 0.4352 |  |
|           | 1988 | 3.04        | 0.09       | 3           | 0.21       |  |
|           | 1989 | 3.18        | 0.11       | 3.38        | 0.28       |  |
| Number of | 1990 | 3.02        | 0.1        | 3.7         | 0.27       |  |
| Diagnosis | 1991 | 3.19        | 0.1        | 3.11        | 0.23       |  |
|           | 1992 | 3.09        | 0.09       | 3.6         | 0.2        |  |
|           | 1993 | 3.42        | 0.11       | 3.65        | 0.22       |  |
|           | 1994 | 3.41        | 0.12       | 3.16        | 0.24       |  |
| Trend     |      | z = -2.8040 | p = 0.0050 | z = -1.1638 | p = 0.2445 |  |
|           | 1988 | 1.52        | 0.06       | 1.44        | 0.09       |  |
|           | 1989 | 1.57        | 0.08       | 1.26        | 0.12       |  |
| Number of | 1990 | 1.6         | 0.04       | 1.73        | 0.16       |  |
| Procedure | 1991 | 1.71        | 0.07       | 1.73        | 0.1        |  |
|           | 1992 | 1.65        | 0.05       | 1.62        | 0.12       |  |
|           | 1993 | 1.69        | 0.05       | 1.59        | 0.18       |  |
|           | 1994 | 1.59        | 0.05       | 1.49        | 0.13       |  |
| Trend     |      | z = 1.5012  | p = 0.1333 | z = 1.4996  | p = 0.1337 |  |

Table6. Distribution of Expected sources of payment (ESOP) by race from 1988 to 1994

|             |      | W           | hite       | African     | American   |  |
|-------------|------|-------------|------------|-------------|------------|--|
| ESOP        | Year | %           | s.e.       | %           | s.e.       |  |
|             | 1988 | 37.8        | 2.57       | 30.3        | 7.4        |  |
|             | 1989 | 40.29       | 2.6        | 45.05       | 6.83       |  |
|             | 1990 | 40.9        | 2.03       | 31.46       | 7.64       |  |
| Medicare    | 1991 | 46.45       | 2.24       | 28.77       | 4.36       |  |
|             | 1992 | 40.94       | 3.08       | 25.86       | 7.1        |  |
|             | 1993 | 42          | 2.43       | 29.05       | 6.65       |  |
|             | 1994 | 38.18       | 2.52       | 23.88       | 4.06       |  |
| Trend       |      | z = 0.3342  | p < 0.7383 | z = -1.9730 | p < 0.0485 |  |
|             | 1988 | 3.35        | 0.94       | 17.66       | 4.45       |  |
|             | 1989 | 3.5         | 0.73       | 13.24       | 7.72       |  |
|             | 1990 | 4.14        | 1.02       | 25.49       | 8.33       |  |
| Medicaid    | 1991 | 4.82        | 1.84       | 19.71       | 6.04       |  |
|             | 1992 | 3.49        | 0.77       | 15.75       | 4.02       |  |
|             | 1993 | 4.97        | 1.06       | 21.41       | 6.3        |  |
|             | 1994 | 4.68        | 1.16       | 27.7        | 6.94       |  |
| Trend       |      | z = 1.1135  | p < 0.2655 | z = 0.8129  | p < 0.4163 |  |
|             | 1988 | 58.85       | 2.71       | 52.04       | 7.92       |  |
|             | 1989 | 56.21       | 2.74       | 41.71       | 6.29       |  |
| Blue cross/ | 1990 | 54.96       | 2.04       | 43.05       | 9.72       |  |
| Other/      | 1991 | 48.73       | 2.29       | 51.53       | 5.69       |  |
| Self        | 1992 | 55.57       | 3.12       | 58.39       | 8.35       |  |
|             | 1993 | 53.03       | 2.66       | 49.54       | 6.23       |  |
|             | 1994 | 57.14       | 2.59       | 48.42       | 6.43       |  |
| Trend       |      | z = -0.7290 | p < 0.4660 | z=0.5281    | p <0.5974  |  |








Figure 2. Hospital admitted breast cancer percentages ( $\pm$  s.e.) of ages 20-44 by race from 1988 to 1994





# Number




Figure 5. Mean number of diagnosis and procedures (95% C.I.) by race from 1988 to 1994






Figure 6. Percentages of routine discharge ( $\pm$  s.e.) by race from 1988 to 1994